



190.30 - Tumor Antigen by Immunoassay CA 19-9

Description

Immunoassay determinations of the serum levels of certain proteins or carbohydrates serve as tumor markers. When elevated, serum concentration of these markers may reflect tumor size and grade. This policy specifically addresses the following tumor antigen: CA19-9.

HCPCS Codes (Alphanumeric, CPT® AMA)

Code	Description
86301	Immunoassay for tumor antigen, quantitative; CA 19-9

ICD-10-CM Codes Covered by Medicare Program

The ICD-10-CM codes in the table below can be viewed on CMS' website as part of Downloads: Lab Code List, at <http://www.cms.gov/Medicare/Coverage/CoverageGenInfo/LabNCDsICD10.html>

Code	Description
C22.1	Intrahepatic bile duct carcinoma
C23	Malignant neoplasm of gallbladder
C24.0	Malignant neoplasm of extrahepatic bile duct
C24.1	Malignant neoplasm of ampulla of Vater
C24.8	Malignant neoplasm of overlapping sites of biliary tract
C24.9	Malignant neoplasm of biliary tract, unspecified
C25.0	Malignant neoplasm of head of pancreas
C25.1	Malignant neoplasm of body of pancreas
C25.2	Malignant neoplasm of tail of pancreas
C25.3	Malignant neoplasm of pancreatic duct
C25.4	Malignant neoplasm of endocrine pancreas
C25.7	Malignant neoplasm of other parts of pancreas
C25.8	Malignant neoplasm of overlapping sites of pancreas
C25.9	Malignant neoplasm of pancreas, unspecified
C78.7	Secondary malignant neoplasm of liver and intrahepatic bile duct
C78.80	Secondary malignant neoplasm of unspecified digestive organ
C78.89	Secondary malignant neoplasm of other digestive organs
D37.6	Neoplasm of uncertain behavior of liver, gallbladder and bile ducts
D37.8	Neoplasm of uncertain behavior of other specified digestive organs



**Medicare National Coverage Determinations (NCD)
Coding Policy Manual and Change Report (ICD-10-CM)**

Code	Description
D37.9	Neoplasm of uncertain behavior of digestive organ, unspecified
G89.3	Neoplasm related pain (acute) (chronic)
M33.03	Juvenile dermatomyositis without myopathy
M33.13	Other dermatomyositis without myopathy
M33.93	Dermatopolymyositis, unspecified without myopathy
R97.8	Other abnormal tumor markers
Z85.068	Personal history of other malignant neoplasm of small intestine
Z85.07	Personal history of malignant neoplasm of pancreas
Z85.09	Personal history of malignant neoplasm of other digestive organs

Indications

Multiple tumor markers are available for monitoring the response of certain malignancies to therapy and assessing whether residual tumor exists post-surgical therapy.

Levels are useful in following the course of patients with established diagnosis of pancreatic and biliary ductal carcinoma. The test is not indicated for diagnosing these two diseases.

Limitations

These services are not covered for the evaluation of patients with signs or symptoms suggestive of malignancy. The service may be ordered at times necessary to assess either the presence of recurrent disease or the patient's response to treatment with subsequent treatment cycles.

ICD-10-CM Codes That Do Not Support Medical Necessity

Any ICD-10-CM code not listed in either of the ICD-10-CM covered or non-covered sections.

Sources of Information

Clinical Pancreatic Guideline for the Use of Tumor Markers in Breast and Colorectal Cancer, American Society of Clinical Oncology. J Clin Oncol 14:2843-2877, 1996.

Richter JM, Christensen MR, Rustgi AK, and Silverstein MD. The Clinical Utility of the CA19-9 Radioimmunoassay for the Diagnosis of Pancreatic Cancer Presenting as Pain or Weight Loss: A Cost Effective Analysis. Arch Intern Med 1989, 149:2292-2297.

Safi F, SchlosseW, Falkenreck S, et. al. Prognostic Value of CA 19-9 Serum Course in Pancreatic Cancer. Hepaetogastroenterology 1998 Jan-Feb; 45(19):253-9.